



Paper Type: Original Article

## Soft Intersection Almost Bi-Quasi Ideals of Semigroups

Aslıhan Sezgin<sup>1,\*</sup> , Aleyna İlgin<sup>2</sup> 

<sup>1</sup>Department of Mathematics and Science Education, Faculty of Education, Amasya University, Amasya, Türkiye; aslihan.sezgin@amasya.edu.tr.

<sup>2</sup>Department of Mathematics, Graduate School of Natural and Applied Sciences, Amasya University, Amasya, Türkiye; aley nailgin@gmail.com.

### Citation:

Received: 24 July 2023

Revised: 09 October 2023

Accepted: 16 January 2024

Sezgin, A., & İlgin, A. (2024). Soft intersection almost bi-quasi ideals of semigroups. *Soft computing fusion with applications*, 1(1), 28-43.

### Abstract


The concept of bi-quasi-ideal generalizes the notions of bi-ideals and quasi-ideals in a semigroup; similarly, the soft intersection bi-quasi-ideal generalizes the concepts of soft intersection bi-ideals and soft intersection quasi-ideals in a semigroup. In this paper, we introduce the concept of soft intersection almost bi-quasi ideal and its generalized concept, soft intersection weakly almost bi-quasi ideals, in a semigroup. In contrast to the soft intersection ideal theory, we demonstrate that every soft intersection almost bi-quasi ideal is also a soft intersection almost ideal and a soft intersection almost bi-ideal. Additionally, we show that every idempotent soft intersection almost bi-quasi ideal is a soft intersection almost subsemigroup, a soft intersection almost weak interior ideal, a soft intersection almost tri-ideal, and a soft intersection almost tri-bi-ideal. Furthermore, we derive several interesting relationships regarding minimality, primeness, semiprimeness, and strong primeness between almost bi-quasi ideals and soft intersection almost bi-quasi ideals with the proven theorem stating that if a nonempty set  $A$  is an almost bi-quasi ideal, then its soft characteristic function is also a soft intersection almost bi-quasi ideal, and vice versa.


**Keywords:** Soft set, Semigroup, Bi-quasi ideals, Soft intersection (Almost) bi-quasi ideals.

## 1 | Introduction

Semigroups are crucial in many branches of mathematics because they provide the abstract algebraic foundation for memoryless systems, which restart on each iteration. Semigroups were first explored formally in the early 1900s. Semigroups are fundamental models for linear time-invariant systems in practical mathematics. Studying finite semigroups is crucial to theoretical computer science, as they are naturally related to finite automata. In addition, semigroups and Markov processes are connected in probability theory.

Ideals are required to comprehend algebraic structures and their uses. In 1952, bi-ideals for semigroups were initially presented by Good and Hughes [1]. The concept of quasi-ideals was introduced by Steinfeld [2] initially for semigroups and then extended to rings. Generalizing ideals in algebraic structures has been a major study area for many mathematicians.

 Corresponding Author: aslihan.sezgin@amasya.edu.tr

 <https://doi.org/10.22105/scfa.v1i1.26>



Licensee System Analytics. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>).

In 1980, Grosek and Satko [3] originally introduced the idea of almost left, right, and two-sided ideals of semigroups. Bogdanovic [4] generalized the notion of bi-ideals to almost bi-ideals in semigroups later, in 1981. In 2018, Wattanatripop et al. [5] defined almost quasi-ideals by combining the ideas of almost ideals and quasi-ideals of semigroups, expanding on the concepts of almost ideals and interior ideals of semigroups, and examining their characteristics. Kaopusek et al. [6] proposed almost interior ideals and weakly almost interior ideals of semigroups in 2020. Iampan et al. [7] in 2021, Chinram and Nakkhasen [8] in 2022, Gaketem [9] in 2022, and Gaketem and Chinram [10] in 2023, Respectively, subsequently introduced almost subsemigroups, almost bi-quasi-interior ideals, almost bi-interior ideals, and almost bi-quasi ideals of semigroups. Additionally, several almost fuzzy semigroup ideal types were investigated in [5], [7]–[12].

In 1999, Molodtsov [13] was the first to propose the idea of the soft set as a way to model uncertainty; this idea has subsequently drawn attention from a variety of fields. In [14]–[23], the fundamental operations of soft sets were examined. Çağman and Enginoğlu [24] modified the idea and presented soft intersection groups [25], which sparked studies on a number of soft algebraic systems. As thoroughly reviewed in [26], [27], soft sets were also conveyed to semigroup theory with the concepts of semigroups with soft intersection left, right, and two-sided ideals, quasi-ideals, interior ideals, and generalized bi-ideals. Different semigroups were categorized by Sezgin and Orbay [28] using soft intersection substructures.

Further research was done on a range of soft algebraic structures in [29]–[38]. Rao [39]–[42] has developed several new semigroup types, including bi-interior ideals, bi-quasi-interior ideals, bi-quasi ideals, quasi-interior ideals, and weak interior ideals, extensions of existing ideals. Furthermore, Baupradist et al. [43] proposed the idea of essential ideals in semigroups.

Rao introduced the bi-quasi ideal of semigroups [41] as a generalization of bi-ideal and quasi-ideal. In contrast, the soft intersection bi-quasi ideal of semigroups was proposed in this paper to generalize the soft intersection bi-ideal and soft intersection quasi-ideal. In [10], almost bi-quasi ideals are introduced as a further generalization of bi-quasi ideals defined in [41]. This study proposes the concept of soft intersection almost bi-quasi ideals and its generalization soft intersection weakly almost bi-quasi ideals of semigroups. Moreover, in contrast to soft intersection semigroup theory, our results show that every soft intersection almost bi-quasi ideal is also a soft intersection almost ideal and a soft intersection almost bi-ideal.

Furthermore, we show that an idempotent soft intersection almost bi-quasi ideal is a soft intersection almost subsemigroup, a soft intersection almost weak interior ideal, a soft intersection almost tri-ideal, and a soft intersection almost tri-bi-ideal. We note that a semigroup may be constructed by soft intersection almost bi-quasi ideals of a semigroup under the binary operation of soft union but not under the soft intersection operation. Also, by deriving that if a nonempty set  $A$  is almost bi-quasi ideal, then its soft characteristic function is also a soft intersection almost bi-quasi ideal, and vice versa, we establish the relationship between a semigroup's soft intersection almost bi-quasi ideal and almost bi-quasi ideal as regards minimality, primeness, semiprimeness, and strongly primeness.

## 2 | Preliminaries

This section reviews several fundamental notions related to semigroups and soft sets.

**Definition 1.** Let  $U$  be the universal set,  $E$  be the parameter set,  $P(U)$  be the power set of  $U$ , and  $K \subseteq E$ . A soft set  $f_K$  over  $U$  is a set-valued function such that  $f_K: E \rightarrow P(U)$  such that for all  $x \notin K$ ,  $f_K(x) = \emptyset$ . A soft set over  $U$  can be represented by the set of ordered pairs [13], [24].

$$f_K = \{(x, f_K(x)): x \in E, f_K(x) \in P(U)\}.$$

Throughout this paper, the set of all the soft sets over  $U$  is designated by  $S_E(U)$ .

**Definition 2.** Let  $f_A \in S_E(U)$ . If  $f_A(x) = \emptyset$  for all  $x \in E$ , then  $f_A$  is called a null soft set and denoted by  $\emptyset_E$ . If  $f_A(x) = U$  for all  $x \in E$ , then  $f_A$  is called an absolute soft set and is denoted by  $U_E$  [24].

**Definition 3.** Let  $f_A, f_B \in S_E(U)$ . If  $f_A(x) \subseteq f_B(x)$  for all  $x \in E$ , then  $f_A$  is a soft subset of  $f_B$  and denoted by  $f_A \subseteq f_B$ . If  $f_A(x) = f_B(x)$  for all  $x \in E$ , then  $f_A$  is called soft equal to  $f_B$  and denoted by  $f_A = f_B$  [24].

**Definition 4.** Let  $f_A, f_B \in S_E(U)$ . The union of  $f_A$  and  $f_B$  is the soft set  $f_A \cup f_B$ , where  $(f_A \cup f_B)(x) = f_A(x) \cup f_B(x)$ , for all  $x \in E$ . The intersection of  $f_A$  and  $f_B$  is the soft set  $f_A \cap f_B$ , where  $(f_A \cap f_B)(x) = f_A(x) \cap f_B(x)$ , for all  $x \in E$  [24].

**Definition 5.** For a soft set  $f_A$ , the support of  $f_A$  is defined by [44]:

$$\text{supp}(f_A) = \{x \in A: f_A(x) \neq \emptyset\}.$$

It is obvious that a soft set with an empty support is a null soft set; otherwise, the soft set is non-null.

**Note 1.** If  $f_A \subseteq f_B$ , then  $\text{supp}(f_A) \subseteq \text{supp}(f_B)$  [45].

A semigroup  $S$  is a nonempty set with an associative binary operation, and throughout this paper,  $S$  stands for a semigroup, and all the soft sets are the elements of  $S_S(U)$  unless otherwise specified.

**Definition 6.** A nonempty subset  $F$  of  $S$  is called:

- I. A left (Right) bi-quasi ideal of  $S$  if  $SF \cap FSF \subseteq F$  ( $FS \cap FSF \subseteq F$ ), and a bi-quasi ideal of  $S$  if  $F$  is both a left bi-quasi ideal of  $S$  and right bi-quasi ideal of  $S$  [41].
- II. An almost left (Right) ideal of  $S$  if  $wF \cap F \neq \emptyset$  ( $Fw \cap F \neq \emptyset$ ), for all  $w \in S$ , and an almost ideal of  $S$  if  $F$  is both an almost left ideal of  $S$  and an almost right ideal of  $S$  [3].
- III. An almost bi-ideal of  $S$  if  $FwF \cap F \neq \emptyset$ , for all  $w \in S$  [4].
- IV. An almost left (Right) bi-quasi ideal of  $S$  if  $(wF \cap Fx) \cap F \neq \emptyset$  ( $(Fw \cap Fx) \cap F \neq \emptyset$ ), for all  $w, x \in S$ , and an almost bi-quasi ideal (Briefly almost BQ-ideal) of  $S$  if  $F$  is both an almost left bi-quasi ideal of  $S$  and an almost right bi-quasi ideal of  $S$  [10].
- V. A weakly almost left (Right) bi-quasi ideal of  $S$  if  $(wF \cap FwF) \cap F \neq \emptyset$  ( $(Fw \cap FwF) \cap F \neq \emptyset$ ), for all  $w \in S$ , and a weakly almost bi-quasi ideal (Briefly weakly almost BQ-ideal) of  $S$  if  $F$  is both a weakly almost left bi-quasi ideal of  $S$  and a weakly almost right bi-quasi ideal of  $S$  [10].

**Definition 7.** An almost left (Right) bi-quasi ideal  $A$  of  $S$  is called a minimal almost left (Right) bi-quasi ideal of  $S$  if for any almost bi-quasi ideal  $B$  of  $S$  if whenever  $B \subseteq A$ , then  $A = B$  [10].

**Definition 8.** Let  $P$  be an almost bi-quasi ideal of  $S$ . Then,  $P$  is called:

- I. A prime almost bi-quasi ideal if for any almost bi-quasi ideals  $A$  and  $B$  of  $S$  such that  $AB \subseteq P$  implies that  $A \subseteq P$  or  $B \subseteq P$ .
- II. A semiprime almost bi-quasi ideal if for any almost bi-quasi ideal  $A$  of  $S$  such that  $AA \subseteq P$  implies that  $A \subseteq P$ .
- III. A strongly prime almost bi-quasi ideal if for any almost bi-quasi ideals  $A$  and  $B$  of  $S$  such that  $AB \cap BA \subseteq P$  implies that  $A \subseteq P$  or  $B \subseteq P$  [10].

**Definition 9.** Let  $f_S$  and  $g_S$  be soft sets over the common universe  $U$ . Then, soft intersection product  $f_S \circ g_S$  is defined by [26].

$$(f_S \circ g_S)(x) = \begin{cases} \bigcup_{x=yz} \{f_S(y) \cap g_S(z)\}, & \text{if } \exists y, z \in S \text{ such that } x = yz, \\ \emptyset, & \text{otherwise.} \end{cases}$$

**Theorem 1.** Let  $b_S, d_S, l_S \in S_S(U)$ . Then,

- I.  $(b_S \circ d_S) \circ l_S = b_S \circ (d_S \circ l_S)$ .
- II.  $b_S \circ d_S \neq d_S \circ b_S$ , generally.
- III.  $b_S \circ (d_S \cup l_S) = (b_S \circ d_S) \cup (b_S \circ l_S)$  and  $(b_S \cup d_S) \circ l_S = (b_S \circ l_S) \cup (d_S \circ l_S)$ .

IV.  $b_S \circ (d_S \tilde{\cap} l_S) = (b_S \circ d_S) \tilde{\cap} (b_S \circ l_S)$  and  $(b_S \tilde{\cap} d_S) \circ l_S = (b_S \circ l_S) \tilde{\cap} (d_S \circ l_S)$ .

V. If  $b_S \tilde{\subseteq} d_S$ , then  $b_S \circ l_S \tilde{\subseteq} d_S \circ l_S$  and  $l_S \circ b_S \tilde{\subseteq} l_S \circ d_S$ .

VI. If  $t_S, k_S \in S_S(U)$  such that  $t_S \tilde{\subseteq} b_S$  and  $k_S \tilde{\subseteq} d_S$ , then  $t_S \circ k_S \tilde{\subseteq} b_S \circ d_S$  [26].

**Definition 10.** Let  $A$  be a subset of  $S$ . We denote by  $S_A$  the soft characteristic function of  $A$  and define as:

$$S_A(x) = \begin{cases} U, & \text{if } x \in A, \\ \emptyset, & \text{if } x \in S \setminus A. \end{cases}$$

The soft characteristic function of  $A$  is a soft set over  $U$ , that is,  $S_A: S \rightarrow P(U)$  [26].

If  $f_S(x) = U$  for all  $x \in S$ , then we denote such a kind of soft set by  $\tilde{\mathbb{S}}$  throughout this paper. It is obvious that  $\tilde{\mathbb{S}} = S_S$ , that is,  $\tilde{\mathbb{S}}(x) = U$  for all  $x \in S$  [26].

**Corollary 1.**  $\text{Supp}(S_A) = A$  [45].

**Theorem 2.** Let  $X$  and  $Y$  be nonempty subsets of  $S$ . Then, the following properties hold [26], [45]:

- I.  $X \subseteq Y$  if and only if  $S_X \tilde{\subseteq} S_Y$ .
- II.  $S_X \tilde{\cap} S_Y = S_{X \cap Y}$  and  $S_X \tilde{\cup} S_Y = S_{X \cup Y}$ .
- III.  $S_X \circ S_Y = S_{XY}$ .

**Definition 11.** Let  $x$  be an element in  $S$ . We denote by  $S_x$  the soft characteristic function of  $x$  and defined as:

$$S_x(y) = \begin{cases} U, & \text{if } y = x, \\ \emptyset, & \text{if } y \neq x. \end{cases}$$

The soft characteristic function of  $x$  is a soft set over  $U$ , that is,  $S_x: S \rightarrow P(U)$  [46].

**Definition 12.** A soft set  $d_S$  of  $S$  over  $U$  is called:

- I. A soft intersection left (Right) bi-quasi ideal of  $S$  over  $U$  if  $(\tilde{\mathbb{S}} \circ d_S) \tilde{\cap} (d_S \circ \tilde{\mathbb{S}}) \tilde{\subseteq} d_S$  ( $(d_S \circ \tilde{\mathbb{S}}) \tilde{\cap} (d_S \circ \tilde{\mathbb{S}}) \tilde{\subseteq} d_S$ ), and a soft intersection bi-quasi ideal of  $S$  if  $d_S$  is both a soft intersection left bi-quasi ideal of  $S$  and a soft intersection right bi-quasi ideal of  $S$ .
- II. A soft intersection almost subsemigroup of  $S$  over  $U$  if  $(d_S \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$  [45].
- III. A soft intersection almost left (Right) ideal of  $S$  over  $U$  if  $(S_x \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$  ( $(d_S \circ S_x) \tilde{\cap} d_S \neq \emptyset_S$ ) for all  $x \in S$ , and a soft intersection almost ideal of  $S$  if  $d_S$  is both a soft intersection almost left ideal of  $S$  and a soft intersection almost right ideal of  $S$  [46].
- IV. A soft intersection almost bi-ideal of  $S$  over  $U$  if  $(d_S \circ S_x \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$  for all  $x \in S$  [47].
- V. A soft intersection almost left (Right) weak interior ideal of  $S$  over  $U$  if  $(S_x \circ d_S \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$  ( $(d_S \circ d_S \circ S_x) \tilde{\cap} d_S \neq \emptyset_S$ ) for all  $x \in S$ , and a soft intersection almost weak interior ideal of  $S$  if  $d_S$  is both a soft intersection almost left weak interior ideal of  $S$  and a soft intersection almost right weak interior ideal of  $S$  [48].
- VI. A soft intersection almost left (Right) tri-ideal of  $S$  over  $U$  if  $(d_S \circ S_x \circ d_S \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$  ( $(d_S \circ d_S \circ S_x \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$ ), for all  $x \in S$ , and a soft intersection almost tri-ideal of  $S$  if  $d_S$  is both a soft intersection almost left tri-ideal of  $S$  and a soft intersection almost right tri-ideal of  $S$  [49].
- VII. A soft intersection almost tri-bi-ideal of  $S$  over  $U$  if  $(d_S \circ d_S \circ S_x \circ d_S \circ d_S) \tilde{\cap} d_S \neq \emptyset_S$  for all  $x \in S$  [50].

It is easy to see that if  $d_S(x) = U$  for all  $x \in S$ , then  $d_S$  is a soft intersection bi-quasi ideal of  $S$ . As is mentioned above, we denote such a kind of soft intersection bi-quasi ideal by  $\tilde{\mathbb{S}}$ . Regarding the probable consequences of network analysis and graph applications concerning soft sets (Defined by the divisibility of determinants), we refer to [51].

### 3 | Soft Intersection Almost Bi-Quasi Ideals of Semigroups

**Definition 13.** A soft set  $f_S$  is called a soft intersection almost left (Right) bi-quasi ideal of  $S$  if

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S \quad ([(f_S \circ S_x) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S),$$

for all  $x, y \in S$ .  $f_S$  is called a soft intersection, the almost bi-quasi ideal of  $S$  if  $f_S$  is both a soft intersection almost left bi-quasi ideal of  $S$  and a soft intersection almost right bi-quasi ideal of  $S$ .

A soft set  $f_S$  is called a soft intersection weakly almost left (Right) bi-quasi ideal of  $S$  if

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_x \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S \quad ([(f_S \circ S_x) \tilde{\cap} (f_S \circ S_x \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S),$$

for all  $x \in S$ .  $f_S$  is called a soft intersection weakly almost bi-quasi ideal of  $S$  if  $f_S$  is both a soft intersection weakly almost left bi-quasi ideal of  $S$  and a soft intersection weakly almost right bi-quasi ideal of  $S$ .

Hereafter, for brevity, a soft intersection is designated by SI, and (Left/right) bi-quasi ideal is designated by (Left/right) BQ-ideal. Thus, soft intersection (Weakly) almost (Left/right) bi-quasi ideal is denoted by SI-(Weakly) almost (Left/right) BQ-ideal.

Here also note that since the operation of soft intersection is commutative in  $S_E(U)$ , it is obvious that in *Definition 13*,  $(S_x \circ f_S)$  and  $(f_S \circ S_y \circ f_S)$  (Similarly,  $(f_S \circ S_x)$  and  $(f_S \circ S_y \circ f_S)$ ) can commute with each other for all  $x, y \in S$ .

**Example 1.** Consider the semigroup  $S = \{m, j\}$  under the binary operation with the following table:

**Table 1. Cayley table of binary operation.**

	$m$	$j$
$m$	$m$	$j$
$j$	$j$	$m$

Let  $f_S$ ,  $h_S$ , and  $g_S$  be soft sets over  $U = D_2 = \{\langle x, y \rangle : x^2 = y^2 = e, xy = yx\} = \{e, x, y, yx\}$  as follows:

$$f_S = \{(m, \{e, x\}), (j, \{e\})\}.$$

$$h_S = \{(m, \{yx\}), (j, \{y, yx\})\}.$$

$$g_S = \{(m, \{x, y\}), (j, \{e, yx\})\}.$$

Here,  $f_S$  and  $h_S$  are both SI-(Weakly) almost BQ-ideal. Let's first show that  $f_S$  is an SI-(Weakly) almost left BQ-ideal, that is,  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ .

Let's start with  $[(S_m \circ f_S) \tilde{\cap} (f_S \circ S_m \circ f_S)] \tilde{\cap} f_S$ :

$$\begin{aligned} & [[(S_m \circ f_S) \tilde{\cap} (f_S \circ S_m \circ f_S)] \tilde{\cap} f_S](m) = [(S_m \circ f_S) \tilde{\cap} (f_S \circ S_m \circ f_S)](m) \cap f_S(m), \\ & = (S_m \circ f_S)(m) \cap (f_S \circ S_m \circ f_S)(m) \cap f_S(m), \\ & = f_S(m) \cap [f_S(m) \cup f_S(j)] \cap f_S(m), \\ & = f_S(m), \end{aligned}$$

$$= f_S(m) \cap f_S(j).$$

Consequently,

$$[(S_m \circ f_S) \tilde{\cap} (f_S \circ S_m \circ f_S)] \tilde{\cap} f_S = \{(m, \{e, x\}), (j, \{e\})\} \neq \emptyset_S.$$

Similarly,

$$[(S_m \circ f_S) \tilde{\cap} (f_S \circ S_j \circ f_S)] \tilde{\cap} f_S = \{(m, \{e\}), (j, \{e\})\} \neq \emptyset_S,$$

$$[(S_j \circ f_S) \tilde{\cap} (f_S \circ S_m \circ f_S)] \tilde{\cap} f_S = \{(m, \{e\}), (j, \{e\})\} \neq \emptyset_S,$$

$$[(S_j \circ f_S) \tilde{\cap} (f_S \circ S_j \circ f_S)] \tilde{\cap} f_S = \{(m, \{e\}), (j, \{e\})\} \neq \emptyset_S.$$

Therefore,  $f_S$  is an SI-(Weakly) almost left BQ-ideal. And also  $f_S$  is an SI-(Weakly) almost right BQ-ideal, that is,  $[(f_S \circ S_x) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . In fact;

$$[(f_S \circ S_m) \tilde{\cap} (f_S \circ S_m \circ f_S)] \tilde{\cap} f_S = \{(m, \{e, x\}), (j, \{e\})\} \neq \emptyset_S,$$

$$[(f_S \circ S_m) \tilde{\cap} (f_S \circ S_j \circ f_S)] \tilde{\cap} f_S = \{(m, \{e\}), (j, \{e\})\} \neq \emptyset_S,$$

$$[(f_S \circ S_j) \tilde{\cap} (f_S \circ S_m \circ f_S)] \tilde{\cap} f_S = \{(m, \{e\}), (j, \{e\})\} \neq \emptyset_S,$$

$$[(f_S \circ S_j) \tilde{\cap} (f_S \circ S_j \circ f_S)] \tilde{\cap} f_S = \{(m, \{e\}), (j, \{e\})\} \neq \emptyset_S.$$

Thus,  $f_S$  is an SI-(Weakly) almost right BQ-ideal. Hence,  $f_S$  is an SI-(Weakly) almost BQ-ideal.

Similarly, we can show that  $h_S$  is an SI-(Weakly) almost left BQ-ideal and SI-(Weakly) almost right BQ-ideal.

Let's first show that  $h_S$  is an SI-(Weakly) almost left BQ-ideal:

$$[(S_m \circ h_S) \tilde{\cap} (h_S \circ S_m \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{yx\})\} \neq \emptyset_S,$$

$$[(S_m \circ h_S) \tilde{\cap} (h_S \circ S_j \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{y, yx\})\} \neq \emptyset_S,$$

$$[(S_j \circ h_S) \tilde{\cap} (h_S \circ S_m \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{yx\})\} \neq \emptyset_S,$$

$$[(S_j \circ h_S) \tilde{\cap} (h_S \circ S_j \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{yx\})\} \neq \emptyset_S.$$

Consequently,  $h_S$  is an SI-(Weakly) almost left BQ-ideal. Let's continue with  $h_S$  is an SI-(Weakly) almost right BQ-ideal:

$$[(h_S \circ S_m) \tilde{\cap} (h_S \circ S_m \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{yx\})\} \neq \emptyset_S,$$

$$[(h_S \circ S_m) \tilde{\cap} (h_S \circ S_j \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{y, yx\})\} \neq \emptyset_S,$$

$$[(h_S \circ S_j) \tilde{\cap} (h_S \circ S_m \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{yx\})\} \neq \emptyset_S,$$

$$[(h_S \circ S_j) \tilde{\cap} (h_S \circ S_j \circ h_S)] \tilde{\cap} h_S = \{(m, \{yx\}), (j, \{yx\})\} \neq \emptyset_S.$$

Therefore,  $h_S$  is an SI-(Weakly) almost right BQ-ideal. Thus,  $h_S$  is an SI-(Weakly) almost BQ-ideal.

One can also show that  $g_S$  is not an SI-(Weakly) almost BQ-ideal. In fact;

$$\begin{aligned}
& [[(S_j \circ g_S) \tilde{\cap} (g_S \circ S_j \circ g_S)] \tilde{\cap} g_S](m) = [(S_j \circ g_S) \tilde{\cap} (g_S \circ S_j \circ g_S)](m) \cap g_S(m), \\
& = (S_j \circ g_S)(m) \cap (g_S \circ S_j \circ g_S)(m) \cap g_S(m), \\
& = g_S(j) \cap [g_S(m) \cap g_S(j)] \cap g_S(m), \\
& = g_S(j) \cap g_S(m), \\
& = \emptyset_S.
\end{aligned}$$

$$\begin{aligned}
& [[(S_j \circ g_S) \tilde{\cap} (g_S \circ S_j \circ g_S)] \tilde{\cap} g_S](j) = [(S_j \circ g_S) \tilde{\cap} (g_S \circ S_j \circ g_S)](j) \cap g_S(j), \\
& = (S_j \circ g_S)(j) \cap (g_S \circ S_j \circ g_S)(j) \cap g_S(j), \\
& = g_S(m) \cap [g_S(m) \cup g_S(j)] \cap g_S(j), \\
& = g_S(m) \cap g_S(j), \\
& = \emptyset_S.
\end{aligned}$$

Consequently,

$$[(S_j \circ g_S) \tilde{\cap} (g_S \circ S_j \circ g_S)] \tilde{\cap} g_S = \{(m, \emptyset), (j, \emptyset)\} = \emptyset_S.$$

Hence,  $g_S$  is not an SI-(Weakly) almost left BQ-ideal. Similarly since

$$[(g_S \circ S_j) \tilde{\cap} (g_S \circ S_j \circ g_S)] \tilde{\cap} g_S = \{(m, \emptyset), (j, \emptyset)\} = \emptyset_S.$$

$g_S$  is not an SI-(Weakly) almost right BQ-ideal and  $g_S$  is not an SI-(Weakly) almost BQ-ideal.

From now on, the proofs are given for only SI-almost left BQ-ideal, since the proofs for SI-almost (Right) BQ-ideal can be shown similarly.

**Proposition 1.** Every SI-almost BQ-ideal is an SI-weakly almost BQ-ideal.

Proof: Let  $f_S$  be an SI-almost BQ-ideal. Then,

$$\begin{aligned}
& [(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S \text{ and } [(f_S \circ S_x) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S \text{ for all } x, y \in S. \text{ Hence,} \\
& [(S_x \circ f_S) \tilde{\cap} (f_S \circ S_x \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S \text{ and } [(f_S \circ S_x) \tilde{\cap} (f_S \circ S_x \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S \text{ for all } x \in S.
\end{aligned}$$

So,  $f_S$  is an SI-weakly almost BQ-ideal.

Since SI-weakly almost BQ-ideal is a generalization of SI-almost BQ-ideal, from now on, all the theorems and proofs are given for SI-almost BQ-ideal instead of SI-weakly almost BQ-ideals.

**Proposition 2.** Let  $f_S$  be an SI-left (Resp. right) BQ-ideal.  $f_S$  is either  $(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) = \emptyset_S$  ( $(f_S \circ S_x) \tilde{\cap} (f_S \circ S_y \circ f_S) = \emptyset_S$ ), for some  $x, y \in S$  or an SI-almost left (Resp. right) BQ-ideal.

Proof: Let  $f_S$  be an SI-left BQ-ideal, then,  $(\tilde{S} \circ f_S) \tilde{\cap} (f_S \circ \tilde{S} \circ f_S) \subseteq f_S$  and let  $(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \neq \emptyset_S$ . We need to show that

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S,$$

for all  $x, y \in S$ . Since  $(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \subseteq (\tilde{S} \circ f_S) \tilde{\cap} (f_S \circ \tilde{S} \circ f_S) \subseteq f_S$ , it follows that  $(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \subseteq f_S$ . From assumption  $(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \neq \emptyset_S$  is obvious. Then,

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S = (S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \neq \emptyset_S.$$

implying that  $f_S$  is an SI-almost left BQ-ideal.

Here it is obvious that, if  $f_S$  is an SI-left BQ-ideal, and for some  $x, y \in S$ ,  $(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) = \emptyset_S$  then,  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S = \emptyset_S \tilde{\cap} f_S = \emptyset_S$ . Therefore,  $f_S$  is not an SI-almost left BQ-ideal.

**Corollary 2.** If  $f_S$  is an SI-almost left (Resp. right) BQ-ideal, then  $f_S$  needs not be an SI-left (Resp. right) BQ-ideal.

**Example 2.** In *Example 1*, it is shown that  $h_S$  is an SI-almost left BQ-ideal; however  $h_S$  is not an SI-left bi-quasi ideal. In fact,

$$\begin{aligned} [(\tilde{S} \circ h_S) \tilde{\cap} (h_S \circ \tilde{S} \circ h_S)](m) &= (\tilde{S} \circ h_S)(m) \cap (h_S \circ \tilde{S} \circ h_S)(m), \\ &= [h_S(m) \cup h_S(j)] \cap [h_S(m) \cup h_S(j)], \\ &= h_S(m) \cup h_S(j), \\ &\not\subseteq h_S(m). \end{aligned}$$

Hence,  $h_S$  is not an SI-left BQ-ideal. Similarly,  $h_S$  is an SI-almost right BQ-ideal; however,  $h_S$  is not an SI-right BQ-ideal. In fact,

$$\begin{aligned} [(h_S \circ \tilde{S}) \tilde{\cap} (h_S \circ \tilde{S} \circ h_S)](j) &= (h_S \circ \tilde{S})(j) \cap (h_S \circ \tilde{S} \circ h_S)(j), \\ &= [h_S(m) \cup h_S(j)] \cap [h_S(m) \cup h_S(j)], \\ &= h_S(m) \cup h_S(j), \\ &\not\subseteq h_S(j). \end{aligned}$$

Hence,  $h_S$  is not an SI-right BQ-ideal.

**Theorem 3.** Every SI-almost left (Resp. right) BQ-ideal is an SI-almost left (Resp. right) ideal.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal. Hence,

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S,$$

for all  $x, y \in S$ . We need to show that  $(S_x \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x \in S$ .

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \cong (S_x \circ f_S) \tilde{\cap} f_S.$$

Since  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(S_x \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Hence,  $f_S$  is an SI-almost left ideal.

**Theorem 4.** Every SI-almost (Left/right) BQ-ideal is an SI-almost bi-ideal.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal. Hence,  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ .

We need to show that  $(f_S \circ S_x \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ .  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \cong (f_S \circ S_x \circ f_S) \tilde{\cap} f_S$

Since  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(f_S \circ S_x \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Hence,  $f_S$  is an SI-almost bi-ideal.

The following example shows that the converse of *Theorem 4* is not true in general:

**Example 3.** Consider the soft set  $g_S$  in *Example 1*. Here,  $g_S$  is an SI-almost bi-ideal, that is,  $(g_S \circ S_x \circ g_S) \tilde{\cap} g_S \neq \emptyset_S$ , for all  $x \in S$ . Let's first show that  $(g_S \circ S_m \circ g_S) \tilde{\cap} g_S \neq \emptyset_S$ :



$$\begin{aligned}
[(g_S \circ S_m \circ g_S) \tilde{\cap} g_S](m) &= (g_S \circ S_m \circ g_S)(m) \cap g_S(m), \\
&= [(g_S(m) \cap (S_m \circ g_S)(m)) \cup (g_S(j) \cap (S_m \circ g_S)(j))] \\
&\cap g_S(m) \\
&= [g_S(m) \cup g_S(j)] \cap g_S(m), \\
&= g_S(m).
\end{aligned}$$

$$\begin{aligned}
[(g_S \circ S_m \circ g_S) \tilde{\cap} g_S](j) &= (g_S \circ S_m \circ g_S)(j) \cap g_S(j), \\
&= [(g_S(m) \cap (S_m \circ g_S)(j)) \cup (g_S(j) \cap (S_m \circ g_S)(m))] \cap g_S(j), \\
&= [g_S(m) \cap g_S(j)] \cap g_S(j), \\
&= g_S(m) \cap g_S(j).
\end{aligned}$$

Consequently,

$$(g_S \circ S_m \circ g_S) \tilde{\cap} g_S = \{(m, \{x, y\}), (j, \emptyset)\} \neq \emptyset_S.$$

Similarly,

$$(g_S \circ S_j \circ g_S) \tilde{\cap} g_S = \{(m, \emptyset), (j, \{e, yx\})\} \neq \emptyset_S.$$

Thus,  $g_S$  is an SI-almost bi-ideal. However, it is clear that  $g_S$  is not an SI-almost (Left/right) BQ-ideal, as seen in *Example 1*.

**Proposition 3.** Let  $f_S$  be an idempotent soft set. If  $f_S$  is an SI-almost (Left/right) BQ-ideal, then  $f_S$  is an SI-almost subsemigroup.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal such that  $f_S$  is an idempotent, then  $f_S \circ f_S = f_S$  and  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . We need to show that  $f_S$  is an SI-almost subsemigroup, that is  $(f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ .

$$\begin{aligned}
[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S &= [(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \tilde{\cap} f_S] \tilde{\cap} f_S \\
&= [(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S) \tilde{\cap} (f_S \circ f_S)] \tilde{\cap} f_S \\
&\cong (f_S \circ f_S) \tilde{\cap} f_S,
\end{aligned}$$

Since  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Thus,  $f_S$  is an SI-almost subsemigroup.

**Proposition 4.** Let  $f_S$  be an idempotent soft set. If  $f_S$  is an SI-almost left (Resp. right) BQ-ideal, then  $f_S$  is an SI-almost left (Resp. right) weak interior ideal.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal such that  $f_S$  is an idempotent, then  $f_S \circ f_S = f_S$  and  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . We need to show that  $f_S$  is an SI-almost left weak interior ideal, that is  $S_x \circ f_S \circ f_S \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x \in S$ .

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \cong (S_x \circ f_S) \tilde{\cap} f_S = (S_x \circ f_S \circ f_S) \tilde{\cap} f_S,$$

Since  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(S_x \circ f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Thus,  $f_S$  is an SI-almost left weak interior ideal.

**Proposition 5.** Let  $f_S$  be an idempotent soft set. If  $f_S$  is an SI-almost left (Right) BQ-ideal, then  $f_S$  is an SI-almost tri-ideal.

Proof: Assume that  $f_S$  is an idempotent SI-almost left BQ-ideal such that  $f_S \circ f_S = f_S$  and  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . We need to show that  $f_S$  is an SI-almost tri-ideal, that is  $(f_S \circ S_x \circ f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$  and  $(f_S \circ f_S \circ S_x \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x \in S$ .

$$[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \subseteq (f_S \circ S_x \circ f_S) \tilde{\cap} f_S = (f_S \circ S_x \circ f_S \circ f_S) \tilde{\cap} f_S.$$

Since  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(f_S \circ S_x \circ f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Hence,  $f_S$  is an SI-almost left tri-ideal.

$$[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \subseteq (f_S \circ S_x \circ f_S) \tilde{\cap} f_S = (f_S \circ f_S \circ S_x \circ f_S) \tilde{\cap} f_S.$$

Since  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(f_S \circ f_S \circ S_x \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Hence,  $f_S$  is an SI-almost right tri-ideal. Thus,  $f_S$  is an SI-almost tri-ideal.

**Proposition 6.** Let  $f_S$  be an idempotent soft set. If  $f_S$  is an SI-almost (Left/right) BQ-ideal, then  $f_S$  is an SI-almost tri-bi-ideal.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal such that  $f_S$  is an idempotent, then  $f_S \circ f_S = f_S$  and  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . We need to show that  $f_S$  is an SI-almost tri-bi-ideal, that is  $(f_S \circ f_S \circ S_x \circ f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x \in S$ .

$$[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \subseteq (f_S \circ S_x \circ f_S) \tilde{\cap} f_S = (f_S \circ f_S \circ S_x \circ f_S \circ f_S) \tilde{\cap} f_S.$$

Since  $[(f_S \circ S_x \circ f_S) \tilde{\cap} (S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $(f_S \circ f_S \circ S_x \circ f_S \circ f_S) \tilde{\cap} f_S \neq \emptyset_S$ . Thus,  $f_S$  is an SI-almost tri-bi-ideal.

**Theorem 5.** Let  $f_S \subseteq h_S$ . If  $f_S$  is an SI-almost left (Resp. right) BQ-ideal, then  $h_S$  is an SI-almost left (Resp. right) BQ-ideal.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal. Hence,  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . We need to show that  $[(S_x \circ h_S) \tilde{\cap} (h_S \circ S_y \circ h_S)] \tilde{\cap} h_S \neq \emptyset_S$ , for all  $x, y \in S$ . In fact,

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \subseteq [(S_x \circ h_S) \tilde{\cap} (h_S \circ S_y \circ h_S)] \tilde{\cap} h_S \neq \emptyset_S.$$

Since  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it is obvious that  $[(S_x \circ h_S) \tilde{\cap} (h_S \circ S_y \circ h_S)] \tilde{\cap} h_S \neq \emptyset_S$ . This completes the proof.

**Theorem 6.** If  $f_S$  and  $h_S$  are SI-almost left (Resp. right) BQ-ideals, then  $f_S \tilde{\cup} h_S$  is an SI-almost left (Resp. right) BQ-ideal.

Proof: Since  $f_S$  is an SI-almost left BQ-ideal and  $f_S \subseteq f_S \tilde{\cup} h_S$ ,  $f_S \tilde{\cup} h_S$  is an SI-almost left BQ-ideal by *Theorem 5*.

**Corollary 3.** The finite union of SI-almost left (Resp. right) BQ-ideals is an SI-almost left (Resp. right) BQ-ideal.

**Corollary 4.** Let  $f_S$  or  $h_S$  be an SI-almost left (Resp. right) BQ-ideal, then  $f_S \tilde{\cup} h_S$  is an SI-almost left (Resp. right) BQ-ideal.

Here, note that if  $f_S$  and  $h_S$  are SI-almost left (Resp. right) BQ-ideals, then  $f_S \tilde{\cap} h_S$  needs not be an SI-almost left (Resp. right) BQ-ideal.

**Example 4.** Consider the SI-almost left (Resp. right) BQ-ideal  $f_S$  and  $h_S$  in *Example 1*. Since,

$$f_S \tilde{\cap} h_S = \{(m, \emptyset), (j, \emptyset)\} = \emptyset_S.$$

$f_S \tilde{\cap} h_S$  is not an SI-almost left (Resp. right) BQ-ideal.

Now, we give the relationship between almost BQ-ideal and SI-almost BQ-ideal. But first of all, we remind the following lemma in order to use it in *Theorem 7*.

**Lemma 1.** Let  $x \in S$  and  $Y$  be a nonempty subset of  $S$ . Then,  $S_x \circ S_Y = S_{xY}$ . If  $X$  is a nonempty subset of  $S$  and  $y \in S$ , then  $S_X \circ S_y = S_{Xy}$  [46].

**Theorem 7.** Let  $A$  be a nonempty subset of  $S$ . Then,  $A$  is an almost left (Resp. right) BQ-ideal if and only if  $S_A$ , the soft characteristic function of  $A$ , is an SI-almost left (Resp. right) BQ-ideal.

Proof: Assume that  $\emptyset \neq A$  is an almost left BQ-ideal. Then,  $(xA \cap AyA) \cap A \neq \emptyset$ , for all  $x, y \in S$ , and so there exist  $j \in S$  such that  $j \in (xA \cap AyA) \cap A$ . Since,

$$\begin{aligned} & \left( [(S_x \circ S_A) \tilde{\cap} (S_A \circ S_y \circ S_A)] \tilde{\cap} S_A \right)(j) = \left( (S_{xA} \tilde{\cap} S_{AyA}) \tilde{\cap} S_A \right)(j), \\ & = \left( (S_{xA \cap AyA}) \tilde{\cap} S_A \right)(j), \\ & = S_{(xA \cap AyA) \cap A}(j), \\ & = U, \\ & \neq \emptyset. \end{aligned}$$

It follows that  $[(S_x \circ S_A) \tilde{\cap} (S_A \circ S_y \circ S_A)] \tilde{\cap} S_A \neq \emptyset_S$ . Thus,  $S_A$  is an SI-almost left BQ-ideal.

Conversely, assume that  $S_A$  is an SI-almost left BQ-ideal. Hence, we have  $[(S_x \circ S_A) \tilde{\cap} (S_A \circ S_y \circ S_A)] \tilde{\cap} S_A \neq \emptyset_S$ , for all  $x, y \in S$ . In order to show that  $A$  is an almost left BQ-ideal, we should prove that  $A \neq \emptyset$  and  $(xA \cap AyA) \cap A \neq \emptyset$ , for all  $x, y \in S$ .  $A \neq \emptyset$  is obvious from the assumption. Now,

$$\begin{aligned} & \emptyset_S \neq [(S_x \circ S_A) \tilde{\cap} (S_A \circ S_y \circ S_A)] \tilde{\cap} S_A, \\ & \Rightarrow \exists \mathcal{S} \in S; \left( [(S_x \circ S_A) \tilde{\cap} (S_A \circ S_y \circ S_A)] \tilde{\cap} S_A \right)(\mathcal{S}) \neq \emptyset, \\ & \Rightarrow \exists \mathcal{S} \in S; \left( (S_{xA} \tilde{\cap} S_{AyA}) \tilde{\cap} S_A \right)(\mathcal{S}) \neq \emptyset, \\ & \Rightarrow \exists \mathcal{S} \in S; (S_{xA \cap AyA} \tilde{\cap} S_A)(\mathcal{S}) \neq \emptyset, \\ & \Rightarrow \exists \mathcal{S} \in S; S_{(xA \cap AyA) \cap A}(\mathcal{S}) = U, \\ & \Rightarrow \mathcal{S} \in (xA \cap AyA) \cap A. \end{aligned}$$

Hence,  $(xA \cap AyA) \cap A \neq \emptyset$ . Consequently,  $A$  is an almost left BQ-ideal.

**Lemma 2.** Let  $f_S \in S_S(U)$ . Then,  $f_S \cong S_{\text{supp}(f_S)}$  [45].

**Theorem 8.** If  $f_S$  is an SI-almost left (Resp. right) BQ-ideal, then  $\text{supp}(f_S)$  is an almost left (Resp. right) BQ-ideal.

Proof: Assume that  $f_S$  is an SI-almost left BQ-ideal. Thus,  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , for all  $x, y \in S$ . In order to show that  $\text{supp}(f_S)$  is an almost left BQ-ideal, by *Theorem 8*, it is enough to show that  $S_{\text{supp}(f_S)}$  is an SI-almost left BQ-ideal. By *Lemma 2*,

$$[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \cong [(S_x \circ S_{\text{supp}(f_S)}) \tilde{\cap} (S_{\text{supp}(f_S)} \circ S_y \circ S_{\text{supp}(f_S)})] \tilde{\cap} S_{\text{supp}(f_S)},$$

and  $[(S_x \circ f_S) \tilde{\cap} (f_S \circ S_y \circ f_S)] \tilde{\cap} f_S \neq \emptyset_S$ , it implies that

$[(S_x \circ S_{\text{supp}(f_S)}) \tilde{\cap} (S_{\text{supp}(f_S)} \circ S_y \circ S_{\text{supp}(f_S)})] \tilde{\cap} S_{\text{supp}(f_S)} \neq \emptyset_S$ . Consequently,  $S_{\text{supp}(f_S)}$  is an SI-almost left BQ-ideal and by *Theorem 7*,  $\text{supp}(f_S)$  is an almost left BQ-ideal.

The following example shows that the converse of *Theorem 8* is not true in general:

**Example 5.** We know that  $g_S$  is not an SI-almost left BQ-ideal in *Example 1*. Since  $\text{supp}(g_S) = \{m, j\}$ ,

$$\begin{aligned} [(m\{m, j\}) \cap (\{m, j\}m\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \\ [(m\{m, j\}) \cap (\{m, j\}j\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \\ [(j\{m, j\}) \cap (\{m, j\}m\{e, j\}\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \\ [(j\{m, j\}) \cap (\{m, j\}j\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \end{aligned}$$

$\text{supp}(g_S)$  is an almost left BQ-ideal. Similarly,  $g_S$  is not an SI-almost left BQ-ideal, but since

$$\begin{aligned} [(\{m, j\}m) \cap (\{m, j\}m\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \\ [(\{m, j\}m) \cap (\{m, j\}j\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \\ [(\{m, j\}j) \cap (\{m, j\}m\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \\ [(\{m, j\}j) \cap (\{m, j\}j\{m, j\})] \cap \{m, j\} &= \{m, j\} \neq \emptyset, \end{aligned}$$

$\text{supp}(g_S)$  is an almost right BQ-ideal. That is to say,  $\text{supp}(g_S)$  is an almost BQ-ideal, although  $g_S$  is not an SI-almost BQ-ideal.

**Definition 14.** Let  $f_S$  and  $h_S$  be SI-almost left (Resp. right) BQ-ideals such that  $h_S \subseteq f_S$ . If  $\text{supp}(h_S) = \text{supp}(f_S)$ , then  $f_S$  is called a minimal SI-almost BQ-ideal.

**Theorem 9.** Let  $A$  be a nonempty subset of  $S$ . Then,  $A$  is a minimal almost left (Resp. right) BQ-ideal if and only if  $S_A$ , the soft characteristic function of  $A$ , is a minimal SI-almost left (Resp. right) BQ-ideal.

Proof: Assume that  $A$  is a minimal, almost left BQ-ideal. Thus,  $A$  is an almost left BQ-ideal, and so  $S_A$  is an SI-almost left BQ-ideal by *Theorem 7*. Let  $f_S$  be an SI-almost left BQ-ideal such that  $f_S \subseteq S_A$ . By *Theorem 8*,  $\text{supp}(f_S)$  is an almost left BQ-ideal and by *Note 1* and *Corollary 1*,

$$\text{supp}(f_S) \subseteq \text{supp}(S_A) = A.$$

Since  $A$  is a minimal almost left BQ-ideal,  $\text{supp}(f_S) = \text{supp}(S_A) = A$ . Thus,  $S_A$  is a minimal SI-almost left BQ-ideal by *Definition 14*.

Conversely, let  $S_A$  be a minimal SI-almost left BQ-ideal. Thus,  $S_A$  is an SI-almost left BQ-ideal, and  $A$  is an almost left BQ-ideal by *Theorem 7*. Let  $B$  be an almost left BQ-ideal such that  $B \subseteq A$ . By *Theorem 7*,  $S_B$  is an SI-almost left BQ-ideal, and by *Theorem 2 (i)*,  $S_B \subseteq S_A$ . Since  $S_A$  is a minimal SI-almost left BQ-ideal,

$$B = \text{supp}(S_B) = \text{supp}(S_A) = A.$$

by *Corollary 1*. Thus,  $A$  is a minimal, almost left BQ-ideal.

**Definition 15.** Let  $f_S$ ,  $g_S$ , and  $h_S$  be any SI-almost left (Resp. right) BQ-ideals. If  $h_S \circ g_S \subseteq f_S$  implies that  $h_S \subseteq f_S$  or  $g_S \subseteq f_S$ , then  $f_S$  is called an SI-prime almost left (Resp. right) BQ-ideal.

**Definition 16.** Let  $f_S$  and  $h_S$  be any SI-almost left (Resp. right) BQ-ideals. If  $h_S \circ h_S \subseteq f_S$  implies that  $h_S \subseteq f_S$ , then  $f_S$  is called an SI-semiprime almost left (Resp. right) BQ-ideal.

**Definition 17.** Let  $f_S$ ,  $g_S$ , and  $h_S$  be any SI-almost left (Resp. right) BQ-ideals. If  $(h_S \circ g_S) \tilde{\cap} (g_S \circ h_S) \subseteq f_S$  implies that  $h_S \subseteq f_S$  or  $g_S \subseteq f_S$ , then  $f_S$  is called an SI-strongly prime almost left (Resp. right) BQ-ideal.

It is obvious that every SI-strongly prime almost left (Resp. right) BQ-ideal is an SI-prime almost left (Resp. right) BQ-ideal, and every SI-prime almost left (Resp. right) BQ-ideal is an SI-semiprime almost left (Resp. right) BQ-ideal.

**Theorem 10.** If  $S_P$ , the soft characteristic function of  $P$ , is an SI-prime almost left (Resp. right) BQ-ideal, then  $P$  is a prime almost left (Resp. right) BQ-ideal, where  $\emptyset \neq P \subseteq S$ .

Proof: Assume that  $S_P$  is an SI-prime almost left BQ-ideal. Thus,  $S_P$  is an SI-almost left BQ-ideal, and thus,  $P$  is an almost left BQ-ideal by *Theorem 7*. Let  $A$  and  $B$  be almost left BQ-ideals such that  $AB \subseteq P$ . Thus, by *Theorem 7*,  $S_A$  and  $S_B$  are SI-almost left BQ-ideals, and by *Theorem 2 (i)* and *(iii)*,

$$S_A \circ S_B = S_{AB} \cong S_P.$$

Since  $S_P$  is an SI-prime almost left BQ-ideal and  $S_A \circ S_B \cong S_P$ , it follows that  $S_A \cong S_P$  or  $S_B \cong S_P$ . Therefore, by *Theorem 2 (i)*,  $A \subseteq P$  or  $B \subseteq P$ . Consequently,  $P$  is a prime almost left BQ-ideal.

**Theorem 11.** If  $S_P$ , the soft characteristic function of  $P$ , is an SI-semiprime almost left (Resp. right) BQ-ideal, then  $P$  is a semiprime almost left (Resp. right) BQ-ideal, where  $\emptyset \neq P \subseteq S$ .

Proof: Assume that  $S_P$  is an SI-semiprime almost left BQ-ideal. Thus,  $S_P$  is an SI-almost left BQ-ideal, and thus,  $P$  is an almost left BQ-ideal by *Theorem 7*. Let  $A$  be an almost left BQ-ideal such that  $AA \subseteq P$ . Thus, by *Theorem 7*,  $S_A$  is an SI-almost left BQ-ideal, and by *Theorem 2 (i)* and *(ii)*,

$$S_A \circ S_A = S_{AA} \cong S_P.$$

Since  $S_P$  is an SI-prime almost left BQ-ideal and  $S_A \circ S_A \cong S_P$ , it follows that  $S_A \cong S_P$ . Therefore, by *Theorem 2 (i)*  $A \subseteq P$ . Consequently,  $P$  is a semiprime almost left BQ-ideal.

**Theorem 12.** If  $S_P$ , the soft characteristic function of  $P$ , is an SI-strongly prime almost left (Resp. right) BQ-ideal, then  $P$  is a strongly prime almost left (Resp. right) BQ-ideal, where  $\emptyset \neq P \subseteq S$ .

Proof: Assume that  $S_P$  is an SI-strongly prime almost left BQ-ideal. Thus,  $S_P$  is an SI-almost left BQ-ideal, and thus,  $P$  is an almost left BQ-ideal by *Theorem 7*. Let  $A$  and  $B$  be almost left BQ-ideals such that  $AB \cap BA \subseteq P$ . Thus, by *Theorem 7*,  $S_A$  and  $S_B$  are SI-almost left BQ-ideals, and by *Theorem 2*,

$$(S_A \circ S_B) \tilde{\cap} (S_B \circ S_A) = S_{AB} \tilde{\cap} S_{BA} = S_{AB \cap BA} \cong S_P,$$

Since  $S_P$  is an SI-strongly prime almost left BQ-ideal and  $(S_A \circ S_B) \tilde{\cap} (S_B \circ S_A) \cong S_P$ , it follows that  $S_A \cong S_P$  or  $S_B \cong S_P$ . Thus, by *Theorem 2 (i)*,  $A \subseteq P$  or  $B \subseteq P$ . Therefore,  $P$  is a strongly prime, almost left BQ-ideal.

## 4 | Conclusion

The concepts of "soft intersection almost bi-quasi ideal" and "soft intersection weakly almost bi-quasi ideal" of semigroups were defined in this work. We demonstrated that although any soft intersection almost bi-quasi ideal is also a soft intersection weakly almost bi-quasi ideal, a soft intersection almost ideal, and a soft intersection almost bi-ideal of a semigroup; the converses are not true in general with counterexamples. Additionally, it was shown that an idempotent soft intersection almost bi-quasi ideal is a soft intersection almost subsemigroup, a soft intersection almost weak interior ideal, a soft intersection almost tri-ideal, and a soft intersection almost tri-bi-ideal. We obtained the relation between soft intersection almost bi-quasi ideal of a semigroup and almost bi-quasi ideal of a semigroup according to minimality, primeness, semiprimeness, and strongly primeness with the obtained theorem that if a nonempty set  $A$  is almost bi-quasi ideal then its soft characteristic function is soft intersection almost bi-quasi ideal, and vice versa. Additionally, we investigated that, unlike soft intersection operation, soft union operation can form a semigroup with the collection of almost bi-quasi ideals of a semigroup. In future studies, some kinds of semigroup ideals, such as quasi-interior ideals, bi-interior ideals, and bi-quasi-interior ideals, may be studied in terms of soft intersection almost ideals.

## Acknowledgment

This paper is derived from the second author's Master Thesis, supervised by the first author at Amasya University, Türkiye.

## Funding

The authors conducted this research independently without external financial support. All costs associated with this study were covered by the researchers' institutional resources.

## Declaration of Interests

The authors affirm that there are no competing interests—financial, professional, or personal—that could influence the objectivity or outcomes of this research. This study was conducted with full academic integrity, free from any affiliations or relationships that might pose a conflict.

## References

- [1] Good, R. A., & Hughes, D. R. (1952). Associated groups for a semigroup. *Bulletin of the American mathematical society*, 58(6), 624–625.
- [2] Steinfeld, O. (1956). Uher die tri ideale, Von halbgruppenn Publ. *Math. debrecen*, 4, 262–275.
- [3] Grošek, O., & Satko, L. (1980). A new notion in the theory of semigroup. *Semigroup forum*, 20(1), 233–240. <https://doi.org/10.1007/BF02572683>
- [4] Bogdanovic, S. (1981). Semigroups in which some bi-ideal is a group. *Review of research faculty of science-university of Novi Sad*, 11, 261–266. [https://sites.dmi.uns.ac.rs/nsjom/Papers/11/NSJOM\\_11\\_261\\_266.pdf](https://sites.dmi.uns.ac.rs/nsjom/Papers/11/NSJOM_11_261_266.pdf)
- [5] Wattanatripop, K., Chinram, R., & Changphas, T. (2018). Quasi-a-ideals and fuzzy a-ideals in semigroups. *Journal of discrete mathematical sciences and cryptography*, 21(5), 1131–1138. <https://doi.org/10.1080/09720529.2018.1468608>
- [6] Kaopusek, N., Kaewnoi, T., & Chinram, R. (2020). On almost interior ideals and weakly almost interior ideals of semigroups. *Journal of discrete mathematical sciences and cryptography*, 23(3), 773–778. <https://doi.org/10.1080/09720529.2019.1696917>
- [7] Iampan, A., Chinram, R., & Petchkaew, P. (2021). A note on almost subsemigroups of semigroups. *International journal of mathematics and computer science*, 16(4), 1623–1629. <https://www.researchgate.net/publication/352899613>
- [8] Chinram, R., & Nakkhasen, W. (2022). Almost bi-quasi-interior ideals and fuzzy almost bi-quasi-interior ideals of semigroups. *Journal of mathematics and computer science*, 26(2), 128–136. <https://doi.org/10.22436/jmcs.026.02.03>
- [9] Gaketem, T. (2022). Almost bi interior ideal in semigroups and their fuzzifications. *European journal of pure and applied mathematics*, 15(1), 281–289. <https://doi.org/10.29020/nybg.ejpam.v15i1.4279>
- [10] Gaketem, T., & Chinram, R. (2023). Almost bi-quasi-ideals and their fuzzifications in semigroups. *Annals of the university of craiova, mathematics and computer science series*, 50(2), 342–352. <https://doi.org/10.52846/ami.v50i2.1708>
- [11] Wattanatripop, K., Chinram, R., & Changphas, T. (2018). Fuzzy almost bi-ideals in semigroups. *International journal of mathematics and computer science*, 13(1), 51–58. <https://www.researchgate.net/publication/322854463>
- [12] Krailoet, W., Simuen, A., Chinram, R., & Petchkaew, P. (2021). A note on fuzzy almost interior ideals in semigroups. *International journal of mathematics and computer science*, 16(2), 803–808. <https://www.researchgate.net/publication/348130224>
- [13] Molodtsov, D. (1999). Soft set theory – first results. *Computers & mathematics with applications*, 37(4-5), 19–31. [https://doi.org/10.1016/S0898-1221\(99\)00056-5](https://doi.org/10.1016/S0898-1221(99)00056-5)
- [14] Ali, M. I., Shabir, M., & Naz, M. (2011). Algebraic structures of soft sets associated with new operations. *Computers and mathematics with applications*, 61(9), 2647–2654. <https://doi.org/10.1016/j.camwa.2011.03.011>
- [15] Sezgin, A., Ahmad, S., & Mehmood, A. (2019). A new operation on soft sets: Extended difference of soft sets. *Journal of new theory*, (27), 33–42. <https://www.researchgate.net/publication/331521192>
- [16] Stojanović, N. S. (2021). A new operation on soft sets: Extended symmetric difference of soft sets. *Vojnotehnički glasnik/military technical courier*, 69(4), 779–791. <https://doi.org/10.5937/vojtehg69-33655>
- [17] Sezgin, A., Aybek, F. N. (2023). A new soft set operation: Complementary soft binary piecewise gamma ( $\gamma$ ) operation. *Matrix science mathematic*, 7(1), 27–45. <https://doi.org/10.54286/ikjm.1304566>
- [18] Sezgin, A., Aybek, F. N., & Atagün, A. O. (2023). A new soft set operation: Complementary soft binary piecewise intersection ( $\cap$ ) operation. *Black sea journal of engineering and science*, 6(4), 330–346. <https://doi.org/10.34248/bsengineering.1319873>

- [19] Sezgin, A., Aybek, F., & Güngör, N. B. (2023). New soft set operation: Complementary soft binary piecewise union operation. *Acta informatica malaysia*, 7(1), 38–53. <http://doi.org/10.26480/aim.01.2023.38.53>
- [20] Sezgin, A., & Demirci, A. M. (2023). A new soft set operation: Complementary soft binary piecewise star (\*) operation. *Ikonion journal of mathematics*, 5(2), 24–52. <http://doi.org/10.54286/ikjm.1304566>
- [21] Sezgin, A., & Yavuz, E. (2023). A new soft set operation: Complementary soft binary piecewise lamda ( $\lambda$ ) operation. *Sinop üniversitesi fen bilimleri dergisi*, 8(2), 101–133. <https://doi.org/10.33484/sinopfb.1320420>
- [22] Sezgin, A., & Yavuz, E. (2023). A new soft set operation: Soft binary piecewise symmetric difference operation. *Necmettin erbakan üniversitesi fen ve mühendislik bilimleri dergisi*, 5(2), 189–208. <https://doi.org/10.47112/neufmbd.2023.18>
- [23] Sezgin, A., & Cagman, N. (2024). A new soft set operation: Complementary soft binary piecewise difference ( $\setminus$ ) operation. *Osmaniye korkut ata üniversitesi fen bilimleri enstitüsü dergisi*, 7(1), 58–94. <https://doi.org/10.47495/okufbed.1308379>
- [24] Çağman, N., & Enginoğlu, S. (2010). Soft set theory and uni-int decision making. *European journal of operational research*, 207(2), 848–855. <https://doi.org/10.1016/j.ejor.2010.05.004>
- [25] Çağman, N., Çitak, F., & Aktaş, H. (2012). Soft int-group and its applications to group theory. *Neural computing and applications*, 21(SUPPL. 1), 151–158. <https://doi.org/10.1007/s00521-011-0752-x>
- [26] Sezer, A. S., Agman, N., Atagün, A. O., Ali, M. I., & Turkmen, E. (2015). Soft intersection semigroups, ideals and bi-ideals; a new application on semigroup theory I. *Filomat*, 29(5), 917–946. <https://doi.org/10.2298/FIL1505917S>
- [27] Sezer, A. S., Çağman, N., & Atagün, A. O. (2014). Soft intersection interior ideals, quasi-ideals and generalized bi-ideals; a new approach to semigroup theory II. *Journal of multiple-valued logic and soft computing*, 23(1–2), 161–207. <https://www.researchgate.net/publication/263651514>
- [28] Sezgin, A., & Orbay, M. (2022). Analysis of semigroups with soft intersection ideals. *Acta universitatis sapientiae, mathematica*, 14(1), 166–210. <https://doi.org/10.2478/ausm-2022-0012>
- [29] Mahmood, T., Rehman, Z. U., & Sezgin, A. (2018). Lattice ordered soft near rings. *Korean journal of mathematics*, 26(3), 503–517. <https://doi.org/10.11568/kjm.2018.26.3.503>
- [30] Jana, C., Pal, M., Karaaslan, F., & Sezgin, A. (2019).  $(\alpha, \beta)$ -soft intersectional rings and ideals with their applications. *New mathematics and natural computation*, 15(2), 333–350. <https://doi.org/10.1142/S1793005719500182>
- [31] Mustuoglu, E., Sezgin, A., & Kaya, Z. (2016). Some characterizations on soft uni-groups and normal soft uni-groups. *International journal of computer applications*, 155(10), 1–8. <https://doi.org/10.5120/ijca2016912412>
- [32] Sezer, A. S., Cagman, N., & Atagün, A. O. (2015). Uni-soft substructures of groups. *Annals of fuzzy mathematics and informatics*, 9(2), 235–246. <https://www.researchgate.net/publication/264792690>
- [33] Sezer, A. S. (2014). Certain characterizations of LA-semigroups by soft sets. *Journal of intelligent and fuzzy systems*, 27(2), 1035–1046. <https://doi.org/10.3233/IFS-131064>
- [34] Özlü, Ş., & Sezgin, A. (2020). Soft covered ideals in semigroups. *Acta universitatis sapientiae, mathematica*, 12(2), 317–346. <https://doi.org/10.2478/ausm-2020-0023>
- [35] Atagun, A. O., & Sezgin, A. (2018). Soft subnear-rings, soft ideals and soft N-subgroups of near-rings. *Mathematical sciences letters*, 7(1), 37–42. <https://doi.org/10.18576/msl/070106>
- [36] Sezgin, A. (2018). A new view on AG-groupoid theory via soft sets for uncertainty modeling. *Filomat*, 32(8), 2995–3030. <https://doi.org/10.2298/FIL1808995S>
- [37] Sezgin, A., Çağman, N., & Atagün, A. O. (2017). A completely new view to soft intersection rings via soft uni-int product. *Applied soft computing journal*, 54, 366–392. <https://doi.org/10.1016/j.asoc.2016.10.004>
- [38] Sezgin, A., Atagün, A. O., Çağman, N., & Demir, H. (2022). On near-rings with soft union ideals and applications. *New mathematics and natural computation*, 18(2), 495–511. <https://doi.org/10.1142/S1793005722500247>
- [39] Rao, M. M. K. (2018). Bi-interior ideals of semigroups. *Discussiones mathematicae-general algebra and applications*, 38(1), 69. <https://doi.org/10.7151/dmgaa.1283>
- [40] Rao, M. M. K. (2018). A study of generalization of bi-ideal, quasi-ideal and interior ideal of semigroup. *Mathematica morovica*, 22(2), 103–115. <http://dx.doi.org/10.5937/MatMor1802103M>

- [41] Rao, M. M. K. (2020). Left bi-quasi ideals of semigroups. *Southeast asian bulletin of mathematics*, 44(3), 369–376. <https://www.researchgate.net/publication/350891820>
- [42] Rao, M. M. K. (2020). Quasi-interior ideals and weak-interior ideals. *Asia pacific journal of mathematics*, 7, 7–21. <https://doi.org/1028924/APIM/7-12>
- [43] Baupradist, S., Chemat, B., Palanivel, K., & Chinram, R. (2021). Essential ideals and essential fuzzy ideals in semigroups. *Journal of discrete mathematical sciences and cryptography*, 24(1), 223–233. <https://doi.org/10.1080/09720529.2020.1816643>
- [44] Feng, F., Jun, Y. B., & Zhao, X. (2008). Soft semirings. *Computers and mathematics with applications*, 56(10), 2621–2628. <https://doi.org/10.1016/j.camwa.2008.05.011>
- [45] Sezgin, A., & İlgin, A. (2024). Soft intersection almost subsemigroups of semigroups. *International journal of mathematics and physics*, 15(1), 13–20. <https://doi.org/10.26577/ijmph.2024v15i1a2>
- [46] Sezgin, A., & İlgin, A. (2024). Soft intersection almost ideals of semigroups. *Journal of innovative engineering and natural science*, 4(2), 466–481. <https://doi.org/10.61112/jiens.1464344>
- [47] Sezgin, A., & Onur, B. (2024). Soft intersection almost bi-ideals of semigroups. *Systemic analytics*, 2(1), 95–105. <https://doi.org/10.31181/sa21202415>
- [48] Sezgin, A., & İlgin, A. (2024). Soft intersection almost weak interior ideals of semigroups :A theoretical study. *Journal of natural and applied sciences Pakistan*, 9(17-18), 372-385. <https://doi.org/10.62792/ut.jnsm.v9.i17-18.p2834>
- [49] Sezgin, A., & İlgin, A. (2024). Soft intersection almost tri-ideals of semigroups. *Journal of innovative engineering and natural science*, 4(2), 466-481. <https://doi.org/10.61112/jiens.1464344>
- [50] Sezgin, A., İlgin, A., & Atagün, A. O. (2024). Soft intersection almost tri-bi-ideals of semigroups. *Science & technology Asia*, 29(4), 1-13. <https://ph02.tci-thaijo.org/index.php/SciTechAsia/article/view/253582>
- [51] Pant, S., Dagtoros, K., Kholil, M. I., & Vivas, A. (2024). Matrices: Peculiar determinant property. *Optimum science journal*, (1), 1–7. <https://www.optimumscience.org/index.php/pub/article/view/9>